

INSECTOSTOP dogs and cats
(external use solution, spot-on)
Package insert

Description

External use solution in a form of transparent oily liquid with specific odour.

Composition

1 ml of the drug contains an active ingredient:

fipronil - 100 mg.

Excipients: propylene carbonate, isopropyl alcohol, citric acid, polyethylene glycol-400.

Pharmacological properties

ATCvet QP53, ectoparasiticides, insecticides and repellents (QP53AX15, fipronil).

Active ingredient of the drug fipronil (Fipronilum)-5-amino-1-[6-dichloro-4-(trifluoromethyl)phenyl]-4-[(trifluoromethyl)sulfinyl]-1H-pyrazole-3-carbonitrile, derivative of phenylpyrazole.

Fipronil hits central nervous system of the ectoparasites, without impact on the mammal's nervous system. Fipronil mode of action in invertebrates is in inhibition of chloride ions passage in GABA-dependent receptors of ectoparasites, disrupting nervous impulses transmission and nervous system activity, including ectoparasites death.

Fipronil is of contact effect and practically not absorbed through the skin. It deposits in epidermis, sebaceous glands of hair follicles and gradually releases for a certain time. It creates an effect of "reservoir" providing fipronil prolonged residual activity.

The drug spreads over the body surface for 24 hours after the topical use.

The drug is toxic for fishes and bees!

Use

Prevention and treatment of cats and dogs with ectoparasite infestations: fleas (*Ctenocephalides* spp.), lice (*Linognathus setosus*) at all stages of development, chewing-lice (*Trichodectes canis*, *Felicola subrostratus*), ixodes ticks (*Ixodes* spp., *Dermacentor* spp., *Rhipicephalus* spp., *Haemaphysalis* spp.).

Treatment of cats and dogs affected with sarcoptic mites (*Otodectes cynotis*, *Sarcoptes scabiei* var. *canis*, *Notoedres cati*, *Cheyletiella* spp.), trombidiformes (*Demodex canis*).

Dosage and administration

The drug is used topical directly on the skin with pipette in the areas inaccessible for animals to lick off: in the area of withers, along the spine, in ear mite infestations – on the skin of auricle.

Drug dosages is the following:

- cats from 3 to 10 kg - 0,8 ml of the drug (1 pipette) per animal;
- dogs:

<i>Dogs body weight, kg</i>	<i>Drug dosage, ml</i>	<i>Number of pipettes, pcs.</i>
3 to 10	0,8	1
10 to 20	1,6	2
20 to 40	2,4	3
over 40	3,2	4

In otodectic mange in animal clean the inner surface of auricle and external auditory channel from ear wax, scabs and exudate, afterwards apply a drug on the skin of both ears, even if the one is affected. Repeat the treatment course in 7-10 days under control of microscopy scrapings in requirement.

In sarcoptic mange, notoedric mange, ear mange and Cheyletiellosis the drug is used 2-4 times in apart 7-10 days. The treatment is recommended to carry out in complex with the use of etiotropic, pathogenetic and symptomatic drugs.

Drug single processing prevents tick infestation within 3 weeks, and fleas up to 4 weeks.

To prevent animals from re-infestation with ectoparasites it is necessary to process beddings with insecticide remedies.

Contraindication

Do not use in animals up to 10-weeks of age!

Do not use the drug in sick, exhausted and debilitated animals!

Do not use in animals with individual hypersensitivity to drug active ingredients!

Use of the drug during the pregnancy and lactation is based on veterinary risk and necessity assessment!

Warnings

It's important to ensure the drug is applied in the areas of the body, which inaccessible for licking

Do not allow the possibility of animals to lick each other after the processing with veterinary drug.

Do not recommended to bath the animals after the processing within 2 days.

The fleas from the pets frequently infect pet's sleeping places and places of stay like carpets and upholstered furniture, thus it's necessary to clean and process with correspondent insecticide remedies.

People with hypersensitivity to insecticides and alcohol do not recommend to process the animals with the drug.

In accident drug getting in the eyes is necessary to rinse carefully with clean water.

Wash the hands with soap thoroughly after the processing of animals with the drug.

Keep the treated animals separated till complete dryness and out of contact with little children.

Release form

Polymeric pipettes of 0,8 ml, 4 or 6 single dose polymeric pipettes in a cardboard packaging

Storage

Keep the drug in closed manufacturer packaging in dry place, out of the rich of children and animals, separate from food and feed at temperature from 4° to 25 °C.

Shelf-life

3 years.

Veterinary use only!

Registration Certificate Holder

“RPE”SUZIRIA” LLC

115, Poltavskii Shlyakh str., Kharkiv, Ukraine 61093

Tel: +38 057 766 36 36

Finished product manufacturer

“RPE”SUZIRIA” LLC

4, Zernova str., Kharkiv, Ukraine 61105

info@provvet.ua

provvet.ua