

INSECTOSTOP collar
(collar)
Package insert

Description

Polyvinyl chloride tape with fastener.

Composition

1,0 g of collar weight (polyvinyl chloride tape) contains active ingredient:
fipronil – 40,0 mg.

Excipients: polyvinyl chloride tape.

Pharmacological properties

ATCvet QP53, ectoparasiticides, insecticides and repellents (QP53AX15, fipronil).

Active ingredient of the drug fipronil (Fipronilum)-5-amino-1-[6-dichloro-4-(trifluoromethyl)phenyl]-4-[(trifluoromethyl)sulfinyl]-1H-pyrazole-3-carbonitrile, derivative of phenylpyrazole.

Fipronil hits central nervous system of the ectoparasites, without impact on the mammal's nervous system. Fipronil mode of action in invertebrates is in inhibition of chloride ions passage in GABA-dependent receptors of ectoparasites, disrupting nervous impulses transmission and nervous system activity, including ectoparasites death.

Fipronil is of contact effect and practically not absorbed through the skin. It deposits in epidermis, sebaceous glands of hair follicles and gradually releases for a certain time. It creates an effect of "reservoir" providing fipronil prolonged residual activity.

The drug is toxic for fishes and bees!

Use

Prevention and treatment of dogs and cats with ectoparasites infestation: fleas (*Ctenocephalides spp.*), lice (*Linognathus setosus*), chewing lice (*Trichodectes canis*, *Felicola subrostratus*), ixodes ticks (*Dermacentor spp.*, *Rhipicephalus spp.*, *Ixodes spp.*, *Haemaphysalis spp.*) at nymph, larvae and imago stages of development.

Dosage and Administration

Unpack the collar and put on the animal neck, fix it loosely in a way to leave a space of 2 cm between the neck and collar, then fasten and cut the excess.

The collar should move freely around the neck to avoid growth restrain of the animal.

Permanent use of the collar protects an animal against ectoparasites within 4 months.

One collar is used per single animal. Use the collar of different length depending on animal size:

- cats, small dog breeds, kittens and puppies of 8-weeks of age – collar of 35 ± 5 cm;

- middle and large dog breeds – collar of 70 ± 5 cm.

It is recommended to process animal beddings and accessories with insecto-acaricide spray INSECTOSTOP.

Contraindications

Do not use in animals up to 8 weeks of age!

Do not use the drug in sick, exhausted and debilitated animals, as well as in animals with intolerance to drug active ingredients!

Do not use in animal within pregnancy and lactation!

Do not use in animals with skin diseases and skin damages!

Warnings

In young animals the collar should be freely on the neck and avoid their growth restrain. Leave a segment of collar while using on kittens and puppies considering their further growth.

Release form

Collar is packed in aluminium foil bag and inserted in cardboard box.

Cats, small dog breeds, puppies and kittens – collar of 35 ±5 cm;
middle and large dog breeds – collar of 70 ±5 cm.

Storage

Keep the drug in closed manufacturer packaging in dry place, out of the rich of children and animals, separate from food and feed at temperature from 4° to 25 °C.

Shelf-life

2 years.

Veterinary use only!

Registration Certificate Holder

“RPE”SUZIRIA” LLC

115, Poltavskii Shlyakh str., Kharkiv, Ukraine 61093

Tel: +38 057 766 36 36

Finished product manufacturer

“RPE”SUZIRIA” LLC

4, Zernova str., Kharkiv, Ukraine 61105

info@provet.ua

provet.ua