

**INSECTOSTOP Spray**  
**(external use solution, spray)**  
Package insert

**Description**

Transparent colorless or light-yellow to yellow color solution with slightly specific odour.

**Composition**

1 ml of the drug contains active ingredient:

fipronil – 3,0 mg.

Excipients: isopropyl alcohol, polyethylene glycol-400.

**Pharmacological properties**

*ATCvet QP53, ectoparasiticides, insecticides and repellents (QP53AX15, fipronil).*

Active ingredient of the drug fipronil (Fipronilum)-5-amino-1-[6-dichloro-4-(trifluoromethyl)phenyl]-4-[(trifluoromethyl)sulfinyl]-1H-pyrazole-3-carbonitrile, derivative of phenylpyrazole.

Fipronil hits central nervous system of the ectoparasites, without impact on the mammal's nervous system. Fipronil mode of action in invertebrates is in inhibition of chloride ions passage in GABA-dependent receptors of ectoparasites, disrupting nervous impulses transmission and nervous system activity, including ectoparasites death.

Fipronil is of contact effect and practically not absorbed through the skin. It deposits in epidermis, sebaceous glands of hair follicles and gradually releases for a certain time. It creates an effect of "reservoir" providing fipronil prolonged residual activity.

The drug spreads over the body surface for 24 hours after the topical use.

The drug is toxic for fishes and bees!

**Use**

Prevention and treatment of dogs and cats with ectoparasites infestation:

fleas (*Ctenocephalides* spp.); lice (*Linognathus setosus*) at all stages of development; chewing lice (*Trichodectes canis*, *Felicola subrostratus*); ixodes ticks (*Dermacentor* spp., *Rhipicephalus* spp., *Ixodes* spp., *Hyalomma* spp., *Haemaphysalis* spp.); acariformes (*Sarcoptes canis*, *Notoedres cati*, *Otodectes cynotis*, *Cheyletiella* spp.).

**Dosage and Administration**

The drug is used external.

Processing of the animals is carried outdoor or in well ventilated premises. The bottle is shaken thoroughly before starting the processing and direct the jet of the drug on the animal body at distance 10-15 cm.

The drug is applied to the entire surface of the animal body, contrary to coat growth, thus to enter not only the coat but on the skin also: at the rate 3-6 ml per 1 kg body weight by pressing the plunger spray – 7 clicks (1ml) for bottles of 100 and 250 ml. It is also recommended to process animal paw pads. In processing the head and areas around the eyes the drug is applied on the hand in gloves and rubbed into the coat, avoiding of getting the drug in mucous membranes.

In otodectic mange in animal clean the inner surface of auricle and external auditory channel from ear wax, scabs and exudate, afterwards apply a drug of 5 ml (3-4 pressings) on the skin of both ears, even if the one is affected. The processing is carried out 2-3 times apart 7-10 days.

In sarcoptic mange, notoedric mange, ear mange and Cheyletiellosis the drug is applied on pre cleaned from the scabs affected areas with taking boundary healthy skin up to 1 cm, at the rate 1 ml (7 pressings) per 10 cm<sup>2</sup> of the affected area.

The processing is carried out 2-4 times apart 7-10 days till clinical recovery of the animal, which confirmed by two negative acarological test results.

In order to prevent animals from ectoparasite infestations is necessary to process their places of stay (dog kennels, cages and etc.) and accessories (beddings, blankets and etc.) at the rate 2 ml (14 pressings) of the drug per 1 m<sup>2</sup>.

The drug is active against ticks up to 30-days and fleas 60-70 days. In processing of long-haired animals the drug dosage is increased in 20-25%. It's not recommended to bath the animal 3 days before and 3 days after the processing.

**Contraindications**

Do not use in animals up to 10 weeks of age!

Do not use the drug in sick, exhausted and debilitated animals!

Do not use in animals with individual hypersensitivity to drug active ingredients!

Use of the drug during the pregnancy and lactation is based on veterinary risk and necessity assessment!

**Warnings**

Pet owners must comply the general hygiene and safety rules adopted for working with veterinary drugs during the processing.

Processing of the animals should be carried outdoor or in well ventilated premises. Do not allow the animals licking the drug.

Avoid of touching the animals after the processing and contact with little children until complete dryness of the drug. In case of getting the drug on the skin or mucous membranes rinse with running water instantly.

Wash the hands with soap thoroughly after the processing of animals with the drug.

**Release form**

Polymeric bottles with sprayer of 100, 250 ml.

**Storage**

Keep the drug in closed manufacturer packaging in dry place, out of the reach of children and animals, separate from food and feed at temperature from 4° to 25 °C.

**Shelf-life**

3 years.

*Veterinary use only!*

**Registration Certificate Holder**

“RPE”SUZIRIA” LLC

115, Poltavskii Shlyakh str., Kharkiv, Ukraine 61093

Tel: +38 057 766 36 36

**Finished product manufacturer**

“RPE”SUZIRIA” LLC

4, Zernova str., Kharkiv, Ukraine 61105

info@provet.ua

provet.ua